



(E) SCAN FOR PRODUCT INFO/SDS







S: ry use. Apply product in short bursts. a to be protected to provide thorough coverage and a titing human activity on the treated area. Repeat applic

DELUTION DIRECTIONS: Disregard when using aerosol can.

Shake or sit EcoBird 14.5 container well prior to diluting. Mk product outside or in

a verillatid eran. Mk EcoBird 14.5 with water at a raiso of 10-50% viv with water. To

are allowed and a supplied and a supplied of the supplied SOFS WY INIXIES, INIX 'pallion' to jloucuc with I spallion to it water.

USE RESTRICTIONS:

Use this product to immediately repel birds from in or on structures, turf areas, landfill and other outdoor areas (including around agricultural buildings and farmsteads). This product must be applied using appropriate spray equipment.

GARBAGE, FOOD STUFFS, WASTE AND TRASH

as warranted by bird activity.

DILUTION DIRECTIONS: Disrepard when using aerosol care.

Shake or sit EcoBird 14.5 container well prior to diluting. For better coverage, add a super weter (such as a non-inior, conganiunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a verifiated area. Mix EcoBird 14.5 with water at a ratio of 10-50% by with water. To make a 20% with mixture, mix 1 gallon of product with 5 gallons of water. To make a 50% with mixture, mix 1 gallon of product with 1 gallon of water.

USE RESTRICTIONS:

Use this product to proper birds from garbage, spilled grain, and other waste food be this product to trape birds from garbage, spilled grain, and other waste food if ir ain is expected.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: EcoBird 14.5 should be stored in its original container in a cool, dry locked place out of reach of children and out of direct sunlight. Do not us or store near heat or open flame.

or store near head or open flame.

PESTICIDE DISPOSAL: To avoid vesates, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a weste disposal facility or pesticide disposal program offers upon program are run by state or local governments or by industry).

CONTAINER DISPOSAL: Norrellible container. Do not reuse or refill this container, imple mase for equivalently promptly after emptying. Empty the remaining contents implement or application of the promption of

WARRANTY STATEMENT

The statements on this label are believed to be true and accurate; however, because usage and conditions which may be or are beyond our control, Roth Chemical Co., and or of the control of the con

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
NI: Harmful if absorbed through sion. Avoid contact with skin, eyes or clot
croughly with soap and water after handing and before eating, drinking,
gun, using tobacco or using the foliet.

wing gum, using tobacco or using the tollet.

ENVIRONMENTAL HAZARDS

terrestrial users, do not apply directly to water, to areas where surface water,
ent, or to intertial asses below the mean high water mark. Do not contamina
er when disposing of equipment washwaters or rinsate.

PERSONAL PROTECTIVE EQUIPMENT (PPE):
idicators and other handlers must wear:
ong-sleeved shift and long paris.

p-sleewed shirt and հարց խառա-ryproof gloves sp plus socks instructions for cleaning/maintaining PPE. Keep and wash PPE tely from other laundry.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its abbeling. Do not apply this product in a way that will contact workers or other perso application. For any requirements specific to your State or Tribe, consult the State or Tribe agency responsible for pesticide regulation.

ribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 GFFR part 170. This standard contains requirements for the protection of agricultural pesticides. It contains requirements for the protection of agricultural pesticides. It contains requirements for their and handlers of agricultural pesticides. It contains requirements for their initial contains and exceptions pertaining to the staetments on this labeling about instructions and exceptions pertaining to the staetments on this labeling about in this box only apply to uses of this product that are covered by the Worker Do not enter or allow worker entry into treated areas during the restricted-entry intervals, and only worker entry into treated areas during the restricted-entry intervals (REI) of zero (0) hours.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticidies (40 CPR Part 170). The WPS applies when this product is used to produce plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas und?

AGRICULTURAL CROPS USE RESTRICTIONS:

Use this product to limit damage caused by birds on ripening crops. This product must be applied using appropriate spray equipment. Do not apply when crop is wet or rain is expected. Withhold treatment until fluid is susceptible to bird damage. Do not apply through any type of irrigation system.

COMPATIBILITY WITH OTHER PESTICIDES:
Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your representative for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the cruly.

residue on the crop.

DILUTION DIRECTIONS:
Shake or stiff EcoBird 14.5 container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. For ground application, mix EcoBird 14.5 with water at a ratio of 1 part product to 10-20 parts water. For aerial application, mix EcoBird 14.5 with water at a ratio of 1 part product to 14-3 parts water.

- oduct to 1-3 parts water.

 PPLICATION DIRECTIONS:
 Apply spray mixture so that crop receives thorough coverage
 Repeat treatment if significant bird damage begins to occur.
 This product may be applied up to the day of harvest.



APPLICATION DIRECTIONS Apply by ground application at a

	of 2.5 – 5 gallons of EcoBird 14.5 (3.1-6.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	
Stone Fruits Apricots Cherries Peaches Plums	Apply by ground application at a rate of 2.5 – 5 gallons of EcoBid 14.5 (3.1-6.2 lbs a.i.) per acre or 7.5-15 ft. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/ handlers and air blast applicators must wear coveralls over normal work clothes and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirts.
Blueberries	Apply by ground application at a rate of 2.5 – 5 gallanos of FcoBild 14.5 (3.1-6.2 lbs. al.) per acre or 7.5-15 ft. cz. per 1000 sgt. ft. ble a 1-10 or 120 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/ handlers and air blast applicators must wear coveralls over normal work clothes and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirts.
Grapes (Table)	Apply by ground application at a rate of 2.5 – 5 gallon of Locality of 1.5 – 5.5 (3.1-8.2 Lbs a.i.) per acre or 7.5-15 ft. (3.1-8.2 Lbs a.i.) per acre or 7.5-15 ft. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers' handlers and air blast applicators must wear coveralis over normal work clothes and not prepare spray solutions using more than 50 gallors of concentrate per day. Harvesters must wear gloves and long-sleeved shirts. New York only: Aerali application is prohibited. This product is not to be used on grapes within Suffolk County.
Cereal Grains Corn (including sweet and pop) Millet Oats	Apply by aerial application at a rate of 2.5 – 5 gallons of EcoBird 14.5 (3.1-6.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:3 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Begin applications when birds begin feeding or crop begins to ripen. New York only: Aerial application is prohibited.
Sunflowers	Apply by aerial application at a rate of 2.5 – 5 gallons of EcoBird 14.5 (3.1-6.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:3 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Begin applications when birds begin feeding or crop begins to ripen. New York only: Aerial application is prohibited.

USE RESTRICTIONS:
Use this product to repel birds from golf courses, other fur areas (including arou agricultural buildings and farmsteada, and hydroseeded areas. This product must be applied using appropriate spray equipment. Do not apply when grass is wet orant is expected.

DILUTION DIRECTIONS:
Shake or set Ecologia 14.5 container well prior to diluting. For better coverage, a super wetter (such as a non-lonic, organofunctional silicone surfactant, or sperader), as directed on the label. Mic product outside or in a ventilated erial EcoBert 14.5 with water at a ratio of 1 part product to 20 parts water. For exam x2 prints of product with 5 gallones of water to make £2.5 gallones of spray mit NP-LCATION DIRECTIONS:

sphy at a rate of 1.2 galants of grary mixture (0.07 lbs. a.i.) per 1000 sq. ft of turf.

sphy at a rate of 1.2 galants of grary mixture (0.07 lbs. a.i.) per 1000 sq. ft of turf.

or by before permitting human activity on the treated area. Repeat application in 4 lays or as warranted by Canada goose activity. Do not mow treated area for severing system of permitting human activity.